

Review for Final Exam 2018/9

I. True or False. ¹

1. The following is valid for all statements p, q, r .

$$\neg(p \wedge q) \vee r \equiv ((\neg p) \vee r) \wedge ((\neg q) \vee r).$$

2. Let A be an $n \times n$ matrix. If a matrix equation $A\mathbf{x} = \mathbf{0}$ has infinitely many solutions, so is $A\mathbf{x} = \mathbf{b}$ for every $n \times 1$ matrix (column vector) \mathbf{b} .

3. Let A be an $m \times n$ matrix with $m < n$. Then a matrix equation $A\mathbf{x} = \mathbf{0}$ has infinitely many solutions.

4. Let $f(x)$ be a function such that $f'(c) = 0$ and $f''(c) = 0$. Then $f(x)$ is either increasing, decreasing or a constant at $x = c$, and $f(c)$ cannot be a local extremum.

5. If $F(x)$ is an antiderivative of $f(x)$, then $F(e^x)$ is an antiderivative of $f(e^x)e^x$.

II. Answer each of the following

1. Write a truth table of $(p \Rightarrow q) \vee ((\neg r) \wedge q)$. ²

2. Find a 3×3 matrix satisfying the following.

$$T \cdot \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a \\ -3a + b \\ a + c \end{bmatrix}$$

3. Find BA and AB . ³

$$A = \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & -1 \\ -2 & 2 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 1 \\ 2 & 1 & 1 \\ -2 & 1 & 2 \end{bmatrix}.$$

4. Determine whether or not the matrix A in the previous problem is invertible. ⁴

5. Find the reduced row echelon form of the matrix below and find the solutions $x_1, x_2, x_3, x_4, x_5, x_6$. ⁵

$$\begin{bmatrix} 1 & 2 & 1 & 0 & -3 & 0 & -2 \\ 0 & 0 & 2 & 0 & 4 & -2 & 0 \\ 0 & 0 & 0 & 1 & 2 & -1 & -5 \\ 0 & 0 & -3 & 0 & -6 & 3 & 0 \end{bmatrix}$$

6. Find a polynomial $f(x)$ of degree three satisfying $f(1) = 1, f(2) = -2, f(3) = 4, f(4) = 12$. ⁶

¹I. FFTFT

²II-1: same as $p \Rightarrow q$, i.e., TTFFTTTT from top in standard order

³II-2: $T = \begin{bmatrix} 1 & 0 & 0 \\ -3 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$, II-3: $AB = \begin{bmatrix} 4 & -1 & -3 \\ 4 & 2 & 1 \\ 4 & 0 & 0 \end{bmatrix}$, $BA = \begin{bmatrix} 0 & 3 & -1 \\ 2 & 3 & -5 \\ -4 & 5 & 3 \end{bmatrix}$

⁴II-4: $A \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 3 \\ 0 & 2 & -4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 3 \\ 0 & 0 & -10 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Thus invertible.

⁵II-5: $\begin{bmatrix} 1 & 2 & 0 & 0 & -5 & 1 & -2 \\ 0 & 0 & 1 & 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 & 2 & -1 & -5 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$, $x_1 = -2s + 5t - u - 2, x_2 = s, x_3 = -2t + u, x_4 = -2t + u - 5, x_5 = t, x_6 = u$.

⁶II-6: $f(x) = \frac{(x-2)(x-3)(x-4)}{(1-2)(1-3)(1-4)} - 2 \frac{(x-1)(x-3)(x-4)}{(2-1)(2-3)(2-4)} + 4 \frac{(x-1)(x-2)(x-4)}{(3-1)(3-2)(3-4)} + 12 \frac{(x-1)(x-2)(x-3)}{(4-1)(4-2)(4-3)}$. II-7. Using $f(x)$ in the previous problem, $f(x) + a(x-1)(x-2)(x-3)(x-4)$, $a \neq 0$ satisfies the condition for any a . Hence there are infinitely many polynomials with the property. II-8. 3.

7. Show that there are infinitely many polynomials $g(x)$ of degree four satisfying $g(1) = 1$, $g(2) = -2$, $g(3) = 4$, $g(4) = 12$.

8. Find the limit $\lim_{n \rightarrow \infty} \frac{3n^3 - 4n + 4}{n^3 - 8}$.

9. Find the limit $\lim_{x \rightarrow 2} \frac{x^2 - 4x + 4}{x^3 - 8}$.

10. Find the limit $\lim_{x \rightarrow 0} \frac{(x+1)e^x - 1}{x}$.

(Hint: Consider the definition of $f'(0)$ when $f(x) = (x+1)e^x$. There is a straight way. You can apply l'Hôpital's rule as well.)

11. Find the derivative of $(3x^2 - 2)^{10}$.

12. Find the derivative of $(x^3 - 2x + 1)e^{-3x^2}$.

13. Find $\int \left(x^2 - \frac{2}{x} + \frac{1}{x^2} \right) dx$.

14. Find $\int_1^4 \left(\sqrt{x} + \frac{1}{\sqrt{x}} \right) dx$.

15. Find $\int x(3x^2 - 2)^9 dx$.

16. Find the derivative of $F(x) = \int_0^x t(3t^2 - 2)^9 dt$.

17. Let $y = f(x) = ce^{ax} + de^{bx}$, where a, b, c, d are constants with $a < b$. Suppose $y'' - 2y' - 3y = 0$ for all c, d , and $f(0) = 2$, $f'(0) = 2$. Find a function $y = f(x)$ with these properties by determining constants a, b, c, d .⁸

III. Answer each of the following

1. Find the inverse of the matrix C below, and solve the system of linear equation.⁹

$$C = \begin{bmatrix} 1 & 0 & -1 & -3 \\ 1 & 0 & -1 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & -2 & 1 & -4 \end{bmatrix}, \quad \begin{cases} x_1 & -x_3 & -3x_4 & = & 1 \\ x_1 & & -x_3 & -2x_4 & = & 2 \\ & x_2 & & +x_4 & = & -3 \\ & -2x_2 & +x_3 & -4x_4 & = & -1 \end{cases}$$

2. Find a function $f(x)$ satisfying $f'(x) = x^2(x-1)(x-5) = x^4 - 6x^3 + 5x^2$ and $f(0) = 1$. Determine whether $f(x)$ has a local maximum, a local minimum, increasing or decreasing at $x = 0, 1, 5$.¹⁰

3. Solve the following differential equations.¹¹

(a) $\frac{dy}{dx} = 3y$, $y(0) = 3$.

(b) $\frac{dy}{dx} = \frac{1}{3\sqrt{xy}}$, $y(1) = 1$, where $x, y > 0$.

⁷II-9:0, II-10:2, II-11: $60x(3x^2-2)^9$, II-12: $(3x^2-2)e^{-3x^2} + (x^3-2x+1)e^{-3x^2}(-6x) = (-6x^4+15x^2-6x-2)e^{-3x^2}$ II-13: $\frac{1}{3}x^3 - 2\log_e|x| - \frac{1}{x} + C$, II-14: $\left[\frac{2}{3}x^{3/2} + 2x^{1/2}\right]_1^4 = \frac{20}{3}$.

⁸II-15:use II-11 to find $\frac{1}{60}(3x^2-2)^{10} + C$, II-16: $x(3x^2-2)^9$, II-17: $y = e^{-x} + e^{3x}$.

⁹III-1: $C^{-1} = \begin{bmatrix} -4 & 5 & 2 & 1 \\ 1 & -1 & 1 & 0 \\ -2 & 2 & 2 & 1 \\ -1 & 1 & 0 & 0 \end{bmatrix}$, $x_1 = -1, x_2 = -4, x_3 = -5, x_4 = 1$.

¹⁰III-2: $f(x) = \frac{1}{5}x^5 - \frac{3}{2}x^4 + \frac{5}{3}x^3 + 1$, increasing at 0, local maximum at 1 and local minimum at 5.

¹¹III-3: (a) $y = 3e^{3x}$. (b) $y = \sqrt[3]{x}$ or $y^3 = x$.